



Miamian Soil Profile

Surface layer: dark grayish brown silt loam Subsoil - upper: dark yellowish brown clay loam Subsoil - lower: yellowish brown clay and clay loam

Substratum: yellowish brown loam

The Miamian series consists of very deep, well drained soils that formed in a thin layer of loess and in the underlying loamy till, which is high in content of lime. The original vegetation consisted of deciduous forest species, principally white oak, maple, elm, ash, and hickory.

Miamian soils are the most extensive soils in Ohio. They occur on more than 750,000 acres in the state. They are productive soils. Corn, soybeans, and winter wheat are the primary crops. The average annual precipitation ranges from 33 to 42 inches, and the average annual air temperature ranges from 51 to 55 degrees F.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

